

PHARMACOGNOSTIC AND THERAPEUTIC ASPECTS OF *ALTERNANTHERA SESSILIS* (L.) IN THE MANAGEMENT OF VARIOUS DISORDERS: A COMPREHENSIVE REVIEW

Twinkle Warkade*, Amol Bondre, Poonam Bihone, Rajesh Mujariya and Manjeet Singh

Institute of Pharmaceutical Science and Research (IPSR), Sardar Patel University, Balaghat (M.P.) India.



*Corresponding Author: Twinkle Warkade

Institute of Pharmaceutical Science and Research (IPSR), Sardar Patel University, Balaghat (M.P.) India.

Article Received on 09/04/2024

Article Revised on 29/04/2024

Article Accepted on 19/05/2024

ABSTRACT

Althernanthera sessilis commonly called as sessile joyweed and dwarf copperleaf is native to tropical and subtropical regions of the world. It is used as a vegetable specially in sri lanka and some asian countries. In certain regions of South East Asia, the leaves and young shoots are consumed as vegetables in karnataka, andhra pradesh and tamil Nadu, the leaves, flowers and tender stems are consumed as vegetables. As an herbal medicine, the plant has diuretic, cooling, tonic and laxative properties. It has been used for the treatment of dysuria and haemorrhoids. The plant is also believed to be beneficial for the eyes, and is used as an ingredient in the making of medicinal hair oils and Kajal.

KEYWORDS: Sessile joyweed, ponnaganti, campesterol, anti-diabetic activity, anti-microbial activity.

INTRODUCTION

Alternanthera sessilis is a prostrate or procumbent, annual or perennial herb. The branches are raised from the root and are up to 50 cm long. They are often purplish in colour and glabrous. The leaves are fleshy, generally 1.3-3 centimetres long and 0.5-1 centimetres wide though the leaves are larger in wet habitats, linear-

elliptic, oblong or obovate, apex rounded and base cuneate. The flowers are inconspicuous, white, borne in small, axillary heads; bracts are ovate or obovate and are 1 mm long. The bracteoles are shorter, persistent; subequal, ovate and acute. The seeds are orbicular. The plant bears flowers and fruits throughout the year.^[1]



Figure 1: *Alternanthera sessilis* shrub.



Figure 2: *A. sessilis* twig with inflorescence.

Alternanthera sessilis Taxonomical Classification^[2]

Domain: Eukaryota	Class: Dicotyledonae
Kingdom: Plantae	Order: Caryophyllales
Phylum: Spermatophyta	Family: Amaranthaceae
Subphylum: Angiospermae	Genus: <i>Alternanthera</i>
	Species: <i>Alternanthera sessilis</i>

Vernacular Names^[3]

Assamese: Matikaduri	Malayalam: Meenamgani, Ponnankannikkira
English: Sessile joyweed	Sanskrit: Matsyaki, Ionica
Hindi: Gudrisag	Kannada: Honagone soppu
Kannada: Honagone soppu	Telugu: Ponnagantikura
Marathi: Bechkusal, Kanchari	Bengali: Chanchi, Haicha, Sachishak
Tamil: Ponnankanni, Citai, Koduppai	Manipuri: Phakchet

Botanical description

A spreading herb, upto 30 cm decumbent herbs, stem glabrous.

Leaves: 3-4 x 1cm, elliptic-oblong to oblanceolate, apex obtuse, base attenuate into a petiole, crenate, glabrous. Spike 7-12 x 5 mm, sessile, solitary or 2-3 together; bracts and bracteoles similar, 1 mm long, broadly ovate, acute, glabrous. This is a perennial herb with prostrate stems, rarely ascending, often rooting at the nodes. Leaves obovate to broadly elliptic, occasionally linear-lanceolate, 1-15 cm long, 0.3-3 cm wide, glabrous to sparsely villous, petioles 1-5 mm long.

Flower: In axillary spikes, sessile; white. Flowering from November-April. Flower in small axillary sessile heads, white often tinged with pink, bracteoles about 1 cm long, ovate, scarious; perianth long, sepals ovate, acute, thin, ovary obcordate, compressed, style very short, capitellate; no characteristic odour and taste. In the wild it flowers from December until March.

Fruit: A cordiform urticle 1.5 mm. long strongly compressed; seed orbicular. orbicular, compressed with thickened margins; no characteristic odour and taste. Fruiting December onwards.

Root: Cylindrical, 0.1 to 0.6 cm. diameter, cream to grey, numerous roots arising from the main tap root as

lateral rootlets; fracture short; no characteristic odour and taste.

Seed: The seed is lenticular, 0.5 to 1 mm in diameter, glossy brown seed coat, slightly reticulate.

Inflorescence: The sessile inflorescences are in axillary position and are isolated or grouped (up to 5). They are subglobose, they measure between 5 and 7 mm in diameter and are white.

Stem: Herbaceous, weak, cylindrical, with spreading branches from the base; yellowish-brown to light-brown; nodes and internodes distinct.^[4,5]

Geographical Distribution

Alternanthera sessilis has a pantropical distribution, being present throughout the Old World tropics, tropical Africa, southern and eastern Asia and Australia. The native range is uncertain (Gupta, 2014). Many Chinese publications (e.g. Fan et al., 2013) describe it as native to China, and USDA-ARS (2014) lists the native range as China and southern to southeastern Asia. It is reported as native to some of the Pacific Ocean island groups and as introduced to others.^[6] However, the most recent study of the genera *Alternanthera* suggests that *A. sessilis* originated in South America and from here was introduced to the Old World.^[7] This is a pioneer species, typically growing on disturbed parts of a variety of

wetland habitats, often in species-rich associations with a range of other aquatic and wetland plants. It grows in the draw- down zones of water bodies or in water up to 1 m deep, where it may be part floating and part emergent or even grow in mats of floating vegetation. It is a typical plant of the flood plain wetlands, margins of rivers, streams, canals, ponds, reservoirs, tanks in India; About 16 species, throughout hotter regions of India and Ceylon in damp places, ascending the Himalayas to 4000 ft, and in all warm countries.^[8,9]

Homotypic Synonyms

<i>Achyranthes sessilis</i> (L.)	<i>Allagathera forsskaolei</i>
<i>Gomphrena sessilis</i> L.	<i>Illecebrum sessile</i> (L.)
<i>Paronychia sessilis</i> (L.)	<i>Paronychia tetragona</i>

Heterotypic Synonyms

<i>Achyranthes linearifolia</i> Sw. ex Wikstr	<i>Achyranthes villosa</i> Blanco
<i>Achyranthes polygonoides</i> (L.)	<i>Alternanthera achyranth</i>
<i>Achyranthes triandra</i> Roxb.	<i>Alternanthera achyranthoides</i>
<i>Achyranthes repens</i> Elliott	<i>Alternanthera denticulata</i> var. <i>bifaria</i>
<i>Alternanthera denticulata</i> var. <i>major</i>	<i>Alternanthera dubia</i>




Figure 3: Pharmacognostic features of *Alternanthera Sessilis*.

Chemical Constituents

Herb contains 2, 4- methylenecycloartanol and cycloeucalenol, choline, oleanolic acid. Saponins have been isolated from the leaves. Roots contain lupeol¹⁷. Young shoots contain protein and iron. It also contains 5- α - stigmaster-7-enol. The β - sitosterol possess potent anti- inflammatory by reducing the secretion of pro inflammatory cytokines and TNF.

Biology and ecology

Alternanthera sessilis prefers places with constant or periodically high humidity, but may however tolerate extremely dry conditions. It often grows in mixed association with several other aquatic species. The propagation of the plant is spreads by seeds, which are wind-and water-dispersed, and by rooting at stem nodes. On basis of Siddha, It grows in Marutham thinai.^[10]

The plant contains an array of chemical constituents viz β sitosterol, stigmasterol, campesterol, α -spinasterol, oleanic acid, rhamnoside, methylene cycloartenol, cycloeucalenol, lupeol, 5- α -stigaster-7-enol and its palmitate, nonacosane, 16-hentriacontane, handianol. oxalic acid, saturated aliphatic hydrocarbon, ester and saturated ester.^[11] Katyakini Muniandy et al., reported that the hydroethanolic extract of *Alternanthera sessilis* showed 50 prominent peaks in GC-MS analysis.




Figure 4: Chemical Structure of β sitosterol & campesterol.

Periyakali saravana Bhavan et al., reported the presence 17 secondary phytochemicals among them 5 compounds possessed bioactive properties.^[12] Kallappa M Hosamani et al., reported the presence ricinolic acid in the seed oil.^[13] Shridhar and Lakshminarayana reported that the leaves of *A. sessilis* contained good amounts α -tocopherol and β -tocopherol.^[14] Ragasa et al., isolated a mixture of diastereomers of new ionone derivatives which showed low anti -microbial activity against *Pseudomonas aeruginosa* and *Trichophyton mentagrophytes*.^[15]

Traditional uses

Alternanthera sessilis (L.) DC. (Sessile joy weed) is a popular leafy vegetable its leaves and young shoots are eaten as vegetable^[16] or cooked in soup in Sri Lanka and also used as traditional medicine in China, Taiwan and India.

Alternanthera sessilis (Amaranthaceae) is widespread throughout the tropics and subtropics and is an annual or perennial prostrate herb with several spreading branches, bearing short petioled simple leaves and small white flowers, found throughout the hotter part of India, ascending to an altitude of 1200m^[17] in the Himalayas and even cultivated as a potherb. The plant grows wild, but is also cultivated for food, herbal medicines, and as an ornamental plant.^[18] The aquarium plant *Alternanthera reineckii* is sometimes misidentified as *A. sessilis*.^[19]

In certain regions of South East Asia, the leaves and young shoots are consumed as vegetables.^[20]

In places like Karnataka, Andhra Pradesh, Tamil Nadu and Sri Lanka the leaves, flowers and tender stems are consumed as vegetables. The plant is shredded finely and stir fried with grated coconut and spices to make a salad-like dish that is most commonly eaten as a rice accompaniment.^[21]

The leaves are crunchy, slightly more so than the temperate climate spinach, and not slimy. Some cultivars are slightly bitter. They require steaming or boiling when eaten in large quantities because of the presence of oxalates.^[22]

It is eaten alone as a green or added to other dishes as a spinach substitute. Reportedly, Brazilians usually eat it raw in salads with oil and or vinegar, tomato, and onion, although the literature recommends cooking it.

The vegetable can be added to quiches, pies, curries, dals, pasta sauces, lasagna or added to dishes and stir-fries late in the cooking process as a spinach substitute and to add a nutty flavour.^[23]

Biochemical analysis were carried out to find total Carbohydrate, Starch, Protein, Amino acid, Vitamin B1 and Vitamin B2 according to the procedure of Association of Official Analytical Chemist. The cooking quality was analyzed for total Carbohydrate, Starch and Proteins. As a herbal medicine, the plant has diuretic, cooling, tonic and laxative properties. It has been used for the treatment of dysuria and haemorrhoids. The plant is also believed to be beneficial for the eyes, and is used as an ingredient in the making of medicinal hair oils and kaja.^[24]

Alternanthera sessilis is known as Matyakshika in Ayurvedic medicine. It has been used in Indian traditional system of medicine since a long time in diseases due to vitiated blood, skin diseases and ulcers.^[25]

Its active principles, extracted in oil, were used to treat infected wounds and the herb also proved styptic in colitis; its nutritive values make the herb a potent tonic with a wide range of applications.

Poultice of pounded fresh material is used in sprains, burns and eczema, carbuncle, erysipelas and acute

conjunctivitis. A decoction is recommended as a herbal remedy to treat wounds, flatulence, nausea, vomiting, cough, bronchitis, diarrhea, dysentery and diabetes. Its roots can relieve inflamed wounds.

The leaves and shoots are boiled and drunk as an antihypertensive remedy. It is also used as a cholagogue, galactagogue, abortifacient, febrifuge and to treat snakebites, dysentery, diarrhea, skin problems inflamed wounds and boils, and applied externally on acne and pimples.

In some parts of Bihar (India) the plant is used for hazy vision, night blindness, and post-natal complaints.

A. sessilis has been reported to possess anti-microbial, molluscicidal, a moderate antimutagenic, antidiarrhoeal, hepatoprotective, cytotoxic and antiviral activities.^[26]

Degenerative and necrotic changes in the liver and kidney in Swiss mice, caused by oral administration of water extract of *A. sessilis* in high doses was revealed through histopathological test.^[27]

Previous phytochemical studies have reported the isolation of flavonols, triterpenoids, steroids and tannins;

β -sitosterol, stigmasterol, campesterol, lupeol being few of its important constituents.

The petroleum ether and benzene extracts inhibit the growth of some human and plant pathogenic bacteria. Previous study on this plant showed that it has hepatoprotective activity and potent nootropic activity.^[28]

The precious method in the Siddha Medicine is 'Food as medicine'. In that way, eating of sessile leaves is fried with ghee, which gives cooling effect to eyes, disease free life. It acts as a good anti-oxidant. Relieves floating abdomen, liver diseases, vatha diseases.

The leaves are very effective in treating in cuts & wounds, antidote for snake bite & scorpion sting and skin diseases. It calcifies copper into red powder (chind), which is very useful in gastric & duodenal ulcers, tamarind should be avoided.

Medicated oil prepared out of the juice of sessile plant as chief ingredient, used as bathing oil, gives cooling effect to eyes & body, neuritis, treating eye, piles, Halitosis. Dried whole plant is used as blood purifier and cures Skin diseases.^[29]

Quantitative microscopy of *Alternanthera sessilis*

S N	Quantitative microscopy of <i>A. sessilis</i>	Numbers
1	Stomatal number	82
2	Stomatal Index	113.8%
3	Vein-islet number	33.5-37/Sq.mm
4	Vein termination number	28-31/Sq.mm
5	Trichome length	1227.4-1431.9 μ m
6	Trichome breadth	122.7-204.6 μ m

CONCLUSION

Our forefathers had a special way of transferring and documenting traditional heritages. one among them is through documenting the traditions through popular sayings for example our focused herb *Alternanthera sessilis* has two such popular sayings, one is About the transformation of body into golden luster and the other one is regarding the clarity of the eyes to visualize the stars even in broad day light when this green is consumed periodically as mentioned in the traditional literatures. These sayings may look like exaggerated once but it has been proved through in-vitro and in-vivo studies that this Green has proven anti-microbial, wound healing activities, anti-oxidant activities, antipyretic activity, nootropic activity, hepatoprotective activity, hematinic activity, anti-ulcer activity, hypoglycemic activity, anti-diarrhoeal property, anti-Inflammatory activity.

REFERENCES

- Kasselmann, Christel Aquarium Plants. Teltow, Germany, 2020; 109. ISBN 978-3-00- 064912-7.
- The Wealth of India, Raw materials, (Revised), CSIR, New Delhi, 1985; 1: 318-9.
- Grubben, G.J.H. & Denton, O.A. Plant Resources of Tropical Africa 2. Vegetables. PROTA Foundation, Wageningen; Backhuys, Leiden; CTA, Wageningen, 2004.
- Holm, L., Doll, J., Holm, E., Pancho, J., Herberger, J. World Weeds: Natural Histories and Distribution. John Wiley & Sons, Inc., New York, USA, 1997.
- Gupta AK, 2014. *Alternanthera sessilis*. The IUCN Red List of Threatened Species. Version, 2014. <http://www.iucnredlist.org>.
- Gupta, A.K, *Alternanthera sessilis*. In: IUCN 2013. IUCN Red List of Threatened Species, 2013. www.iucnredlist.org.
- The Siddha Pharmacopoeia of India, Publication by Ministry of Health & Family Welfare, Department of AYUSH, 2.
- Gupta A.K. Indian medicinal plants New Delhi. ICMAR, 2004; 151-7.
- Sharma PC, Yelne MB, Dennis TJ, Database on medicinal plants used in Ayurveda, central council of research in Ayurveda and siddha, New delhi, 14: 398.
- Periyakali BS, Kanagarasu R Rajkumar G, Virumandi N, TangaveluS, Thangaraj M,

- Phytochemical characterization of *Alternanthera sessilis* and Assessment of its growth promoting potential on the fresh water prawn *Macrobrachium rosenbergii*, International journal of research studies in Zoology, 2017; 3(4): 25-38.
11. Hosamani KM, Ganjihal SS Chavadi DV, *Alternanthera triandra* seed oil: a moderate source of ricinoleic acid and its possible utilization, Industrial crops and products, 2004; 19(2): 133-136.
 12. Shridhar R, Gollamudi LN, Lipid classes, fatty acids and tocopherol of leaves of six edible plants species, *Journal of Agric. Food. chem.*, 1993; 41: 61-63.
 13. Ragasa CY, Tremor N, Redeont JA, Ionone derivatives from *Alternanthera sessilis*, J. Asian. Nat. prod. Res., 2002; 4(2): 109-115.
 14. Barukial J, Sarmah JN, Ethnomedicinal plants used by the people of Goleghat district, Assam, *International journal of medicinal and aromatic plants*, 1(3): 203-211.
 15. Reddy KN, Pattanaik C, Reddy CS, Raju VS, Traditional knowledge on wild food plants in Andrapradesh, Indian journal of traditional knowledge Jan., 2007; 6(1): 223-229.
 16. Mathur M, Nama KS, Choudhary K, Economic and ethno medicinal importance of floral diversity on ancient walls of Kota district, Rajasthan, International journal of pure and applied biosciences, 2016; 4(4): 167-173.
 17. Jiji P, Ethno medicinal uses of wild vegetables used by Tai-shyam people of Sivasagar district, Assam, India, International Research journal of biological sciences, Nov., 2014; 3(11): 63-65.
 18. Panda A, Mishra MK, Ethno medicinal survey of some wetland plants of south Orissa and their conservation, Indian journal of traditional knowledge April, 2011; 10(2): 296-303.
 19. Horo S, Topno S, Study and analysis of nutritional value of some wild and semi wild edible plants by "HO" tribes of Singhbhum district, Jharkhand, India, 2015; 3(5): 25-32.
 20. Hiremath VT, Vijaykumar MMJ, Taranath TC, Survey on ethno medicinal plants of jogi matti forest, chitradurga district, Karnatka, India, Environment & We an international journal of science and technology, 2010; 5: 223-233.
 21. Jenny MOL PA, Suganthi A, Ethnobotanical survey on medicinal plants used by the tribal people in Attappady, Kerala, International journal of pharmaceutical sciences and research, Jan., 2017; 2(1): 17-23.
 22. Singhal A, Khare RK, Yadav R, Comparative study of some ethno medicinal plants among the tribals of Datia and Sheopurkalan, District (M.P), International journal of life sciences scientific research Jan., 2017; 838-843.
 23. Vinod kumar S, Gadepa S, Koochamgari N, Babu rao N, An ethnobotanical survey of medicinal plants used by ethnic people in Bodhan, Nizamabad district, Telangana state, India, IOSR journal of pharmacy, Sep., 2017; 7(9): 53-5.
 24. Sharma P, Joshi A, Some important medicinal plants of Dehradun zoo, Uttarkhand, world journal of pharmacy and pharmaceutical sciences, 2018; 7(40): 1099-1103.
 25. Bharat B Maitreya, Ethnomedicinal values of some weed plant species of Bhavnagar, Gujarat, India, International journal of emerging trends in science and technology, June, 2015; 2(6): 2675-2681.
 26. Tiwari AP, Joshi B, Ansari AA, Ethno medicinal uses of some weeds of Uttarpradesh, India, Researcher, 2012; 4(7).
 27. Prasad SK, Singh BN, Documentation of ethno medicinal plants of Gopalganj district of Bihar (India) IOSR Journal of pharmacy and biological sciences, version -3, May-June, 2014; 9(3): 80-89.
 28. Savita rani and Rana JC, Ethno botanical uses of some plants of Bhattiyat block in district chamba. Himachal Pradesh (Western Himalaya), Ethno botany research and applications, 2014; 12: 407-414.
 29. Shiddamallayya N, Yasmeen A, Gopalkumar K, Hundred common forest medicinal plants of Karnataka in primary health care, Indian journal of traditional knowledge Jan., 2010; 9(1): 90-95. Accessed on 2.5.2019.